

ALLEGAN COUNTY HEALTH DEPARTMENT

3255 – 122ND Ave., Suite 200, Allegan, MI 49010

Office Administration

(269) 673-5411

Bioterrorism Preparedness

673-5411

Personal Health

673-5411



Communicable Disease

673-5411

Environmental Health

673-5415

Resource Recovery

673-5415

MEASLES

What is measles?

Measles is a respiratory illness with rash caused by the rubeola virus.

Who can get measles?

Anyone who has not received 2 documented doses of MMR or has not had a confirmed case of measles can get measles. If exposed, approximately 90% of people who have not been vaccinated or previously had measles will develop the disease.

What are the symptoms of measles?

Measles starts with a fever and at least one of the "three C"s: **c**ough, **c**oryza, and **c**onjunctivitis. It's followed by a rash that starts on the face and spreads over the body.

- High fever (over 101° F)
- Cough
- Coryza (runny nose)
- Conjunctivitis (red, watery eyes)
- Tiny white spots on the inner cheeks, gums, and roof of the mouth (Koplik Spots) 2-3 days after symptoms begin
- A rash that is red, raised, blotchy; usually starts on face, spreads to trunk, arms, and legs 3-5 days after symptoms begin

How long after exposure do symptoms begin?

Symptoms usually begin 7-14 days after exposure, but can appear as long as 21 days after exposure. If symptoms develop, call ahead before you visit your doctor, urgent care or emergency room so they can take precautions to prevent exposure to other individuals.

How is measles spread?

Measles is a very contagious disease that is easily spread by person-to-person direct contact and airborne spread of droplets from the nose, throat, and mouth through sneezing, coughing, and speaking. It is spread through even minimal air exposure and contact with contaminated surfaces.

How long is a person contagious?

Measles can be spread 4 days before developing the rash through 4 days after the rash appears. Measles is highly contagious. A person can be infected with measles just by being in a room with an infected person, even up to 2 hours after the infected person has left.

I have been exposed to someone with measles, what do I do?

- **Unvaccinated individuals need to get vaccinated within 72 hours of exposure.** If you do not have a record of receiving two measles (MMR) vaccines, unsure if you have been vaccinated, or unsure if you have had measles in the past, contact your healthcare provider.
- **Immune Globulin (IG) treatment is effective within 6 days of exposure for high-risk individuals** including those who are unvaccinated or unsure about vaccination status, pregnant women, and those with a weakened immune system due to illness and diseases like HIV, malnutrition, and/or medications.
- **If symptoms develop, call ahead before you visit your doctor or emergency room** so they can take precautions to prevent exposure to other individuals.
- **Stay home if you are sick and don't allow visitors in your home as measles is highly contagious.**
- **Watch for symptoms for 21 days after potential exposure.** Call your healthcare provider if symptoms develop and you believe you were exposed.

Are there complications from measles?

Measles can make individuals very ill. Encephalitis (inflammation of the brain) is a serious complication of measles. A more common complication of measles is an ear infection or pneumonia. During pregnancy, measles may cause preterm birth or the baby being born at a low birthweight. Death due to measles is rare in the U.S.

Is there a treatment for measles?

If exposed to someone with measles, unvaccinated individuals or those without evidence of immunity can be vaccinated within 72 hours of exposure to help prevent disease. Immune Globulin (IG) treatment is effective within 6 days of exposure for high-risk individuals.

Bed rest at home for at least 4 days after the appearance of the rash is necessary to not infect others. Supportive care can be given to relieve symptoms and address complications.

How can measles be prevented?

- Measles is preventable by vaccination.
- If you do not have a record of two documented measles (MMR) vaccines from a doctor or in the Michigan Care Improvement Registry (MCIR), unsure if you were vaccinated, or unsure if you had measles in the past, contact your healthcare provider about getting vaccinated.
- Measles vaccine is a 2-dose series. If you have two documented doses of the MMR vaccine you do not need another dose.
- All healthy children should be vaccinated at 12 - 15 months with the combination shot for measles, mumps, rubella (MMR).
- A 2nd MMR vaccine is usually given at 4 - 6 years of age, but can be given earlier if it is at least 4 weeks after the 1st dose. Children who receive two MMR vaccines at least 4 weeks apart at any time after the age of 12 months are considered to have completed the MMR series.
- Some infants under 12 months should get a dose of MMR if they are in an area with ongoing transmission or are traveling out of the country. This dose will not count toward their routine series, so these children would still need to receive two MMR vaccines after 12 months of age. Contact your healthcare provider to determine if your infant should get a dose of MMR.

- Adults born in 1957 or later should receive at least 1 dose of MMR vaccine unless they have other acceptable evidence of immunity. A second dose of MMR vaccine is needed for adults who may have been exposed to a measles case, or those who are students in colleges/universities, work in health care, or plan to travel internationally. Check with your health care provider to see if you need to be vaccinated.
- In an outbreak situation: potentially exposed individuals without documentation of 2 MMR doses should receive a dose of MMR.
- Pregnant women should not get vaccinated as this is a live viral vaccine.
- All women of childbearing age should avoid contact with those who have measles.
- Be sure to keep a record of all immunizations. Write down when the shots were received.

What if measles occurs at school or a day care center?

All cases must be reported to your local health department within 24 hours. People born in 1957 or later who cannot prove that they either have had: *Laboratory evidence of immunity to measles, or Measles vaccine after 12 months of age and a 2nd dose at least 4 weeks later*, should get a measles vaccine. Otherwise, they will be excluded from school/day care for at least 21 days after the beginning of the last measles case.

More information can be found at www.cdc.gov or www.michigan.gov/measlesoutbreak

Please contact your healthcare provider if you have concerns regarding your or your child's health. If you have general questions, you may contact an Allegan County Public Health Nurse at 269-673-5411.